Pharmaceutical Analysis Notes PDF | B Pharm 1st Semester Notes

---

Unit 1(a): Introduction to Pharmaceutical Analysis

Pharmaceutical analysis is like being a detective for medicines. It's a crucial branch of practical chemistry that helps us understand everything about a given substance or sample. This involves breaking it down, separating its parts, figuring out what it is (identification), how much of it is there (quantification), and making sure it's super clean (purification).

What do pharmaceutical analysts do? Their work is vital in many areas, including:

- **Checking raw materials**: Ensuring that the initial ingredients used to make drugs are of good quality.
- **Analysing drug samples**: Testing various medicines to confirm their composition.
- **Qualitative and quantitative analysis**: Discovering *what* is in a sample (qualitative) and *how much* is present (quantitative).
- **Diagnosing diseases**: Using chemical analysis to help understand illnesses.
- **Working with radioactive compounds**: Determining the properties of these special substances.
- **Ensuring water quality**: Analysing water samples to make sure they are safe.
- **Quality control**: Making sure drugs are effective, safe, and pure.
- **Forensic science**: Aiding in investigations.
- **Plant growth**: Finding out the amount of essential components in plants.

In essence, qualitative analysis tells you **"what it is"**, identifying the components or functional groups of a substance. Quantitative analysis answers **"how much"**, determining the precise amount of a substance in a sample.

Techniques in Pharmaceutical Analysis

To solve the mysteries of pharmaceutical compounds, analysts use various techniques. These are broadly categorised:

1. **Chemical Methods**: These are the traditional hands-on approaches.

    - **Volumetric Methods (Titrimetric Analysis)**: Imagine measuring out exactly how much liquid you need to react with another liquid. That's volumetric analysis! The key is measuring the volume of a solution (called the **titrant**) of known strength that reacts completely with the substance being analysed (the **analyte** or **titrate**). The amount of analyte is then calculated from the volume of titrant used.
        - **Titration**: This is the process where you slowly add the titrant from a **burette** into the analyte, which is usually in a **conical flask**, until the reaction is complete.
        - **Equivalence Point** (or Stoichiometric End Point): This is the theoretical point where the reaction between the titrant and analyte is perfectly complete.
        - **Indicator**: To see when the equivalence point is reached, we often add a special substance called an **indicator**. It changes colour visibly at or near the equivalence point, which we call the **end point**.
        - **Titration Error**: Sometimes, the practical end point (where the indicator changes colour) might be slightly different from the theoretical equivalence point. This difference is the titration error.
        - **Types of Volumetric Methods**: These include neutralisation titrations (acid-base), precipitation titrations, complexometric titrations, non-aqueous titrations, and oxidation-reduction titrations.
        - **Important considerations for volumetric methods**: The reaction should be easy, complete, without interference, fast, and compatible with suitable indicators that give sharp colour changes.
    - **Gravimetric Methods**: Instead of volume, this method focuses on **weight**. You isolate the compound you're interested in, usually in its pure form, and weigh it. This is particularly useful for checking impurities but can be quite time-consuming. Examples include methods based on precipitation or volatilisation.
    - **Gasometric Methods**: As the name suggests, these methods involve the measurement of gases.

2. **Physico-chemical Methods (Instrumental Methods)**: These are modern methods that use sophisticated instruments to measure physical or physicochemical properties of compounds.
    - Examples of properties measured and the instruments used:
        - **Electrical potential**: Potentiometer
        - **Electrical conductance**: Conductometry
        - **Electrical current**: Polarography and Voltammetry
        - **Absorption of radiation**: Spectrophotometry, Colorimetry, Atomic absorption spectroscopy
        - **Emission of radiation**: Emission spectroscopy, Flame photometry, Fluorimetry
        - **Scattering of radiation**: Turbidimetry and Nephelometry
        - **Refraction of radiation**: Refractometry
        - **Rotation of plane polarized light**: Polarimetry, Optical rotatory dispersion
        - **Thermal properties**: Thermal methods of analysis (e.g., DSC, DTA, TGA)
        - **Mass to charge ratio**: Mass spectrometry
    - Other instrumental techniques include HPLC, HPTLC, Gas Chromatography, and TLC.
3. **Microbiological Methods**: These methods are used when chemical tests aren't suitable, especially for compounds like antibiotics. They involve observing how a substance affects the **growth of bacteria** compared to a standard.
4. **Biological Methods**: These are used to determine the concentration of a drug and to check its overall activity or effect.

Methods of Expressing Concentrations

When we talk about solutions in pharmaceutical analysis, we need to be precise about how much solute (the substance dissolved) is in the solvent (the dissolving liquid). Here are the common ways to express concentration:

- **Normality (N)**: This is the number of **equivalents** of solute present in one litre of solution. It's often calculated as: Normality (N) = Weight of Solute / (Equivalent Weight of Solute × Volume of Solution in Litres) It is also known as Normal Concentration.
- **Percent Concentration (%)**: This expresses the amount of solute as parts per hundred. There are three main types:
    - **Percent w/w (% w/w)**: Mass of Solute / Mass of Solution × 100. Used for commercial aqueous reagents.
    - **Percent v/v (% v/v)**: Volume of Solute / Volume of Solution × 100. Used when diluting one liquid with another.
    - **Percent w/v (% w/v)**: Mass of Solute / Volume of Solution × 100. Used when dissolving solid reagents in solvents.
- **Molar Concentration (Molarity, M)**: This is the number of **moles** of solute present in one litre of solution. It's calculated as: Molarity (M) = Number of Moles of Solute / Volume of Solution in Litres Molarity can change slightly with temperature because volume changes.
- **Molal Concentration (Molality, m)**: This is the number of **moles** of solute per 1000 grams (or 1 kilogram) of solvent. Unlike molarity, molality is not affected by temperature as it depends on mass, not volume.
- **Formal Concentration (Formality, F)**: For substances that exist as ions in solution, like salts, we use formality. Instead of molecular weight, we use **formula weight**. Formality (F) = Weight of Solute in Grams / (Formula Weight × Volume of Solution in Litres) Like molarity, formality can be influenced by temperature.
- **Parts per Million (ppm) and Parts per Billion (ppb)**: These units are used for very dilute solutions, especially when dealing with impurities. For example, 1 ppm means 1 part of solute in one million parts of solution. Concentration (ppm) = Mass of Solute / Mass of Solution × 10^6 ppm

Equivalent Weight The equivalent weight of a substance is the weight that reacts with specific amounts of other substances (like 1.0078 g of hydrogen or 8 g of oxygen). Its calculation depends on the type of reaction:

- **Neutralisation Reactions (Acids/Bases)**:
    - For acids: Equivalent weight = Molecular weight / Basicity (number of replaceable hydrogens).
    - For bases: Equivalent weight = Molecular weight / Acidity (number of replaceable hydroxyl groups).

- **Complexation Reactions**: For cations, it's generally the atomic weight divided by the valency of the atom.
- **Precipitation Reactions**: Equivalent weight = Molecular weight / Valency of reacting ions.
- **Oxidation-Reduction Reactions**: Here, the equivalent weight is based on the number of electrons transferred or the change in oxidation number. Equivalent weight = Molecular weight / Number of electrons transferred Or, Equivalent weight = Molecular weight / Change in oxidation number
    - **Oxidation Number**: A positive change means oxidation, a negative change means reduction. Some general rules for determining oxidation numbers exist: non-zero digits are significant, zeros between non-zero digits are significant, trailing zeros after a decimal point are significant, leading zeros are never significant, exact numbers have infinite significant digits.

Titer: This term refers to the concentration of a solution, particularly in volumetric analysis. It tells you the weight of a substance that will react with the solute in 1 ml of the solution. Titer = Normality (N) × Equivalent Weight

---

Preparation and Standardization of Various Molar and Normal Solutions

In pharmaceutical analysis, having solutions of precisely known concentrations is fundamental. These are called **standard solutions**.

- **Primary Standards**: These are the gold standard! They are substances that are:
    - Easily available in a highly pure and crystalline form.
    - Used to prepare standard solutions of unknown strength through accurate weighing.
    - **Properties**: They react simply and completely, have no difference between their theoretical and practical end points, don't absorb water from the air (not hygroscopic), are readily soluble in common solvents, easy to purify and preserve, react immediately and completely, are stable in the air, and have a high equivalent weight (which minimises weighing errors).
    - Examples include oxalic acid and sodium carbonate.

- **Secondary Standards**: These are solutions whose concentrations are determined by comparing them against a primary standard. They are used because primary standards are limited or might not be suitable for all reactions.
    - **Properties**: Their reactions might not be as simple or instantaneous, they might be difficult to keep pure, their concentration can change over time (e.g., due to decomposition or absorbing water), and they might not be readily soluble.
    - An example is potassium permanganate ($KMnO_4$), which is hygroscopic.

Let's look at how some common solutions are prepared and standardised:

- **Preparation and Standardisation of N/10 Sodium Hydroxide (NaOH) Solution**:

    - **Preparation**: Carefully weigh about 4 grams of NaOH, dissolve it in distilled water, transfer to a 1-litre volumetric flask, cool, and then fill to the mark with distilled water. Mix well.
    - **Standardisation**: Titrate this NaOH solution against a known N/10 oxalic acid solution using phenolphthalein as an indicator. The end point is a faint permanent pale pink colour. You can then calculate the exact normality of your NaOH solution using the formula $N_1V_1 = N_2V_2$.

- **Preparation and Standardisation of N/10 Sulphuric Acid (H_2SO_4) Solution**:

    - **Preparation**: Since concentrated sulphuric acid is very strong, you need to calculate the exact volume required (e.g., about 2.74 ml of 97% concentrated H_2SO_4 for 1 litre of N/10 solution). Carefully add this calculated volume to a small amount of distilled water in a beaker, then transfer to a 1-litre volumetric flask and make up to the mark with distilled water. Mix well.
    - **Standardisation**: Titrate this H_2SO_4 solution against a standard 0.1 N sodium carbonate (Na_2CO_3) solution, using mixed indicator or methyl orange. Repeat until you get consistent readings and use $N_1V_1 = N_2V_2$ for calculation.

- **Preparation of N/10 Oxalic Acid Solution**:

    - **Preparation**: Accurately weigh 6.3 grams of oxalic acid dihydrate ($(COOH)_2.2H_2O$) and transfer it into a 1-litre volumetric flask. Fill halfway with distilled water, shake

well to dissolve, and then make up to the mark. This solution is often used as a primary standard. (If anhydrous oxalic acid is available, 4.5 g is used for 0.1 N).

- **Preparation and Standardisation of N/10 Potassium Permanganate ($KMnO_4$) Solution**:
    - **Preparation**: The equivalent weight of $KMnO_4$ depends on the reaction. In an acidic medium (most common), it's 31.6 (158/5). So, weigh approximately 3.16 to 3.2 grams of $KMnO_4$, dissolve it in 1 litre of distilled water. Boil the solution for 10-15 minutes, let it stand for a few days, and then filter through glass wool to remove any impurities.
    - **Standardisation**: Take 10 ml of N/10 oxalic acid solution, add 5 ml of dilute sulphuric acid, and warm it to 60-70°C. Titrate this warm solution against the $KMnO_4$ from the burette until a light pink colour appears ($KMnO_4$ acts as its own indicator). Repeat for consistent results and use $N_1V_1 = N_2V_2$ for calculation.

- **Preparation and Standardisation of N/10 Hydrochloric Acid (HCl) Solution**:
    - **Preparation**: Calculate the exact volume of concentrated HCl needed (e.g., about 8.1774 ml of 37.5% concentrated HCl for 1 litre of 0.1 N solution). Dissolve this volume in 1 litre of water.
    - **Standardisation**: Titrate this HCl solution against a standard N/10 NaOH solution (which has already been standardised against oxalic acid) using phenolphthalein as an indicator.

- **Preparation and Standardisation of 0.1 N Sodium Thiosulphate ($Na_2S_2O_3.5H_2O$) Solution**:
    - **Preparation**: Dissolve approximately 24.8 grams of sodium thiosulphate crystals in previously boiled and cooled distilled water and make the volume up to 1 litre. Store the solution in a cool, dark-coloured bottle to prevent decomposition.
    - **Standardisation**: Weigh about 5.0 grams of finely ground potassium dichromate (a primary standard) into a 1-litre volumetric flask and make it up to volume. Pipette 25.0 ml of this dichromate solution into a conical flask. Add 5.0 ml of concentrated hydrochloric acid and 15.0 ml of 10% potassium iodide solution. Allow it to stand in the dark for 5 minutes. Then titrate the mixture with your sodium thiosulphate

solution, using starch solution as an indicator. The end point is when the blue colour changes to green.

Unit 1(b): Pharmaceutical Errors

In analytical chemistry, getting the correct result every time is the goal. However, sometimes errors happen. An **error** is simply the difference between the **true result** (the accepted correct value) and the **measured result** you obtain. If these errors are large, they can have serious consequences, especially in medicine, leading to expensive or even dangerous treatments for patients.

Understanding Errors:

- **Error Calculation**: Error = Measured Mean Value - True Value
- **Absolute Error**: This is the straightforward difference between your experimental average and the true value.
- **Relative Error**: This error is more about proportion; it's the absolute error divided by the true value. It's often expressed as a percentage (by multiplying by 100) or as parts per thousand (by multiplying by 1000).

Sources and Types of Errors: Errors in pharmaceutical analysis come from various sources and are generally classified into two main types:

1. **Determinate Errors (Systematic Errors)**: These errors are **known to the analyst** and, importantly, **can be avoided or corrected**. They tend to be constant or reproducible errors.

    - **Instrumental Errors**: Caused by faulty equipment, uncalibrated instruments, or instruments that don't perform correctly.
    - **Personal Errors**: These happen because of the analyst themselves. This could be due to a lack of experience, carelessness, poor observation skills (like vision problems), or improper sampling.
    - **Chemical Errors**: Result from using chemicals or reagents that have impurities or contaminants, which interfere with the reaction.

    - **Errors in Methodology**: These are very serious errors arising from using a flawed or incorrect method. Examples include impurities precipitating along with the desired substance (**co-precipitation**), substances not dissolving properly, or incomplete reactions. These are usually detectable and can often be eliminated.
    - **Additive Errors**: Errors that are not dependent on the amount of substance present in the sample.
    - **Proportional Errors**: Errors that depend on the amount and nature of the sample.
2. **Indeterminate Errors (Accidental or Random Errors)**: These errors **may or may not be known** and are often **unpredictable**. They show up as small, random differences in a series of measurements taken under identical conditions.

    - You **cannot predict or eliminate** these errors. They follow a random pattern, which can be understood using probability.
    - Generally, very large random errors are unlikely, while smaller errors happen more frequently. These errors can occur on both the positive and negative sides of the true value with equal probability.

Methods to Minimise Errors: Since errors can't always be completely eliminated, we focus on minimising them:

- **Personal Errors**: Employ skilled and knowledgeable individuals for analysis. Implement regular reporting and monitoring. Check for issues like vision problems.
- **Instrumental Errors**: Always thoroughly check and **calibrate** all equipment before starting any analysis. Faulty equipment should be corrected by experts.
- **Chemical Errors**: Use only high-quality, **standard chemicals from authentic sources** that are free from impurities. Regularly check the quality of reagents.
- **Errors in Methodology**: Strictly follow **standard methods** and proper references. Continuous monitoring of reactions by experienced personnel helps.
- **Indeterminate Errors**: While unpredictable, a **well-planned analytical procedure** that considers all factors influencing accuracy and precision can help reduce their impact.
- **Calibration of Instruments/Apparatus**: This is the process of checking the correctness of your measuring tools.
- **Blank Determination**: Perform the analysis without the sample to identify any impurities in the reagents or solvents themselves.

- **Control Determination**: Conduct the analysis with known quantities of the substance to detect errors.
- **Independent Methods**: Use two different analytical methods to measure the same substance and compare the results.
- **Parallel Determination**: Repeat the analysis more than once (e.g., in duplicate or triplicate) and compare the results to ensure consistency.

Accuracy and Precision

These two terms are often used interchangeably, but in chemistry, they have distinct meanings:

- **Accuracy**: This refers to the **degree of closeness of a measurement to the true or target value**. Think of it as hitting the bullseye on a dartboard. If your measurements are accurate, they are very close to what they *should* be. Accuracy is sometimes called **bias error**. High accuracy requires both high precision and high "trueness" (how close the average of many measurements is to the true value).
- **Precision**: This refers to the **degree of agreement between several measurements of the same quantity**. It's about **reproducibility**. If your measurements are precise, they are very close to each other, even if they aren't necessarily close to the true value. Imagine hitting the same spot on the dartboard repeatedly, but that spot isn't the bullseye.

Good accuracy and good precision means all your darts are clustered around the bullseye. **Poor accuracy but good precision** means all your darts are tightly clustered, but far from the bullseye. **Poor accuracy and poor precision** means your darts are scattered all over the board.

Significant Figures

When recording measurements in analysis, it's important to use **significant figures** correctly. These are the number of digits that are necessary to express the result of a measurement in a way that is consistent with its precision. They tell you which part of your measurement is certain and which part has a bit of uncertainty (usually the last digit).

Rules for Identifying Significant Figures:

- **Non-zero digits are always significant**. (e.g., 123 has three significant figures).
- **Captive zeros** (zeros between two non-zero digits) are **always significant**. (e.g., 25.05 has four significant figures).
- **Trailing zeros** (zeros at the end of a number) are **significant only if there is a decimal point** in the number. (e.g., 0.0560 has three significant figures, because of the decimal. 1000 without a decimal has one significant figure, but 1000. has four).
- **Leading zeros** (zeros before non-zero digits) are **never significant**. They just show the magnitude or placement of the decimal point. (e.g., 0.0234 has three significant figures).
- **Exact numbers** (like counts or definitions) are considered to have an **infinite number of significant digits**.

Rounding Off Figures: Sometimes, in calculations or when reporting data, you need to round off numbers to maintain the correct number of significant figures for accuracy.

- In addition and subtraction, the result should have the same number of decimal places as the measurement with the fewest decimal places.
- In multiplication and division, the result should have the same number of significant figures as the measurement with the fewest significant figures.

Unit 1(c): Pharmacopoeia, Sources of Impurities & Limit Test

Pharmacopoeia

Imagine a giant rulebook for medicines – that's essentially a **Pharmacopoeia**! The word comes from Greek: "pharmakon" (drug or medicine) and "pieo" (to make). These are official publications that provide a comprehensive list of drugs and medicinal substances, along with detailed instructions for their preparation and quality control.

- **What's in a Pharmacopoeia?**
    - **Quality Specifications**: It sets out appropriate tests to confirm the identity, purity, amount of active substance, and performance characteristics of medicines.
    - **Reference Standards**: It specifies reference standards that are used for testing to ensure consistent quality.

    - **Coverage**: It covers everything from raw materials, inactive ingredients (excipients), and intermediate products to the final finished pharmaceutical products (FPPs).
    - **General Requirements**: It also includes analytical methods, microbiological purity tests, dissolution testing, stability guidelines, and more.
- **International Pharmacopoeia (Ph. Int.)**: Issued by the World Health Organization (WHO), this provides global standards.
- **Key Function**: Pharmacopoeias are essential tools for maintaining the **quality** of medicines by providing recommended procedures for analysis and specifications for pharmaceutical substances, excipients, and dosage forms.
- **Structure**: Most Pharmacopoeias have a **general part** (tests, methods, general requirements) and a **specific part** consisting of **monographs** for individual pharmaceutical substances.

The Indian Pharmacopoeia (IP): In India, drugs manufactured must be labelled with the mandatory "I.P." (Indian Pharmacopoeia).

- **Publisher**: The Indian Pharmacopoeia is published by the **Indian Pharmacopoeial Commission (IPC)**, under the Government of India, Ministry of Health & Family Welfare. It adheres to the requirements of the Drugs and Cosmetics Act, 1940.
- **Mission**: The IPC's mission is to **promote public health** by ensuring the quality, safety, and effectiveness of medicines, including active pharmaceutical ingredients, excipients, and dosage forms.
- **History**: The first IP was published in 1868. Over the years, it has been updated regularly, with new editions and addenda incorporating scientific advancements and focusing on national health care priorities. For example, the **Indian Pharmacopoeia 2022** introduced 92 new monographs, 27 new APIs, vitamins, minerals, amino acids, fatty acids, biotechnology products, herbal products, blood products, dosage forms, and vaccines.
- **Objectives of IPC**:
    - To develop and update **comprehensive monographs** for drugs.
    - To create monographs specifically for **herbal drugs**.
    - To prioritise monographs for drugs on the **National Essential Medicines List**.
    - To consider the latest analytical testing and instrumentation when framing monographs.
    - To accelerate the process of preparing, certifying, and distributing IP Reference Substances.

    - To collaborate with other international pharmacopoeias (like Ph Eur, BP, USP) to harmonise global standards.
    - To review existing monographs periodically.
    - To organise educational programs and research.
    - To publish the National Formulary of India.
- **General Chapters**: The IP includes general chapters covering notices, test methods (biological, chemical, physical, pharmaceutical), reference data, reagents, general tests, containers, and tables.

Monographs: A **monograph** in a Pharmacopoeia is like a detailed dossier for an individual official substance or preparation.

- **General Monographs**: These provide requirements for various dosage forms (e.g., tablets, capsules) and set general rules for their production and manufacture.
    - **Production**: Contains mandatory instructions for manufacturing processes, covering raw materials, in-process testing, storage, and validation.
    - **Manufacture of Drug Products**: Focuses on the active ingredient, ensuring all other ingredients (excipients) also meet Pharmacopoeial requirements.
    - **Excipients**: These are inactive substances added to medicines. Monographs ensure they are harmless, don't affect the drug's effectiveness, and don't interfere with tests.
- **Individual Monographs**: These provide specific details for individual drug products. They include:
    - **Titles**: The official name, often the International Non-proprietary Name (INN).
    - **Chemical Formulae**: The chemical structure, molecular formula, and molecular weight, referring to the chemically pure substance.
    - **Definition**: The official definition of the drug product.
    - **Statement of Content**: Limits of the active ingredient determined by assay.
    - **Category**: Indicates its medical or pharmaceutical basis.
    - **Dose and Usual Strength**: Information for pharmacists and medical practitioners.
    - **Description**: A preliminary evaluation of the preparation.
    - **Solubility**: Information on approximate solubility.

Sources of Impurities

Even in highly pure substances, some amount of **impurity** is almost always present. An impurity is any foreign particle or unwanted chemical that affects the purity of a substance. In pharmaceuticals, these are unwanted chemicals that remain with the **active pharmaceutical ingredients (APIs)** or develop during the formulation process. Even tiny amounts can impact the effectiveness and safety of medicines.

Where do impurities come from?

- **Raw Materials**: The initial materials used in manufacturing might not be entirely pure.
- **Manufacturing Process**: The methods or processes used to make the drug can introduce impurities.
- **Chemical Reactions**: The chemical processes themselves, or the plant materials used, can lead to unwanted by-products.
- **Storage Conditions**: Improper storage (e.g., exposure to light, air, or moisture) can cause degradation and form impurities.
- **Decomposition**: The drug substance itself can break down over time, creating impurities.

What are the effects of impurities?

- **Incompatibility**: Impurities can react negatively with other substances in the formulation.
- **Reduced Shelf Life**: They can make the medicine expire faster.
- **Formulation Difficulties**: Impurities can complicate the process of making the drug.
- **Altered Properties**: They might change the physical and chemical properties of the drug.
- **Decreased Therapeutic Effect**: The medicine might not work as well as it should.
- **Toxic Effects**: Impurities can be harmful or toxic, especially if present above certain limits.
- **Sensory Changes**: They can change the odour, colour, or taste of the drug.

How can we minimise pharmaceutical impurities?

- **Purity of Starting Materials**: Always use very pure starting ingredients.
- **Optimised Processes**: Minimise impurity levels during synthesis and manufacturing.
- **Impurity Identification**: Identify the chemical structures of impurities.
- **Toxicity Studies**: Isolate and synthesise impurities to evaluate their potential toxicity.
- **Stability Monitoring**: Constantly monitor the stability of APIs and drug products (DPs) to detect any degradation.
- **Proper Storage**: Ensure correct storage and stability analysis.

Tests for Purity: Pharmacopoeias include specific "tests for purity" to detect impurities and set limits for how much impurity is acceptable.

- **Insoluble Residue**: Pure substances dissolve clearly. Cloudiness or opalescence indicates insoluble impurities.
- **Ash, Water Insoluble Ash**: This test checks for inorganic impurities like heavy metals or minerals, particularly in crude plant-derived drugs.
- **Colour, Odour, and Taste**: While limited, these sensory characteristics can give a preliminary indication of purity.
- **Physico-chemical Constants**: Properties like solubility, melting and boiling points, optical rotation, and refractive index can help assess a substance's purity.
- **Acidity, Alkalinity, and pH**: Tests for these properties help estimate impurities arising from acids or alkalis used in chemical reactions.
- **Anions and Cations**: Common impurities like chloride and sulphate ions often result from the use of strong acids (hydrochloric, sulphuric, nitric) during synthesis.

Limit Test

Limit tests are crucial quantitative tests designed to detect and control small quantities of specific impurities present in a substance. They are based on comparing the turbidity (cloudiness) or colour produced in a sample with that produced in a standard solution.

Why are Limit Tests important?

- To identify harmful amounts of impurities.
- To determine both avoidable and unavoidable impurities.

Let's explore some specific limit tests:

- **Limit Test for Chloride**:
    - **Principle**: This test relies on the reaction between silver nitrate and soluble chloride ions in an acidic environment (dilute nitric acid) to form a cloudy precipitate of silver chloride (opalescence). The opalescence produced in your sample is then compared to a standard solution. If your sample shows less opalescence than the standard, it passes the test.
    - **Chemical Reaction**: Cl- + AgNO3 --(Dilute HNO3)--> AgCl (precipitate) + NO3-

    - **Observation**: The sample solution's opalescence should not be greater than that of the standard solution.
- **Limit Test for Sulphate**:
    - **Principle**: This test involves the reaction of soluble sulphate ions with barium chloride in the presence of dilute hydrochloric acid to form insoluble barium sulphate, which appears as a turbidity. This turbidity is compared to a standard. The barium sulphate reagent also contains alcohol, which prevents excessive supersaturation and helps in forming a uniform turbidity.
    - **Chemical Reaction**: SO4^2- + BaCl2 --(HCl)--> BaSO4 (precipitate) + 2Cl-
    - **Reasons**: Hydrochloric acid makes the solution acidic, potassium sulphate enhances sensitivity, and alcohol helps prevent supersaturation.
    - **Observation**: The sample solution's opalescence should not be greater than that of the standard solution.
- **Limit Test for Iron**:
    - **Principle**: Iron in an ammoniacal (alkaline) solution, in the presence of citric acid and thioglycollic acid, forms an iron thioglycolate complex, which is pale pink to deep reddish-purple. Thioglycollic acid reduces ferric iron to ferrous iron. Citric acid prevents the precipitation of iron by ammonia, keeping it in solution. The colour intensity of the sample is compared with a standard.
    - **Reasons**: Citric acid forms a soluble complex with iron, preventing its precipitation by ammonia. Thioglycollic acid acts as a reducing agent (Fe(III) to Fe(II)) and then forms the coloured complex. Ammonia makes the solution alkaline, which is necessary for the colour development.
    - **Observation**: The purple colour produced in the sample solution should not be greater than the standard solution.
- **Limit Test for Heavy Metals**:
    - **Principle**: This test is based on the reaction of metallic impurities with hydrogen sulphide in an acidic medium, forming coloured sulphides (typically reddish or blackish). The amount of metallic impurities is expressed as parts of lead per million (ppm), with the Indian Pharmacopoeia usually setting a limit of 20 ppm. Metals that respond to this test include lead, mercury, bismuth, arsenic, antimony, tin, cadmium, silver, copper, and molybdenum.
    - **Chemical Reaction**: Heavy Metal + H2S --> Sulphide of Heavy Metal (brownish colour) + 2H+

    - **Observation**: The colour produced in the sample solution should not be greater than the standard solution.
- **Limit Test for Lead**:
    - **Principle**: This test involves a chemical reaction between lead and diphenyl thiocarbazone (Dithizone) in an alkaline solution to form a violet-coloured lead-dithizone complex. Dithizone itself is green, so the resulting solution turns red. Reagents like ammonium citrate, potassium cyanide (KCN), and hydroxylamine hydrochloride ($NH_2OH.HCl$) are used to prevent interference from other metals and to optimise the pH.
    - **Observation**: The intensity of the red colour in the sample solution should not be greater than that of the standard solution.

- **Limit Test for Arsenic (Gutzeit Test)**:
    - **Principle**: This test, known as the Gutzeit Test, involves converting all arsenic present into arsine gas (AsH_3) by reducing it with zinc and hydrochloric acid. The arsine gas then reacts with mercuric chloride paper, producing a yellow stain. The intensity of this yellow stain is proportional to the amount of arsenic present and is compared against a standard stain. The IP typically sets a limit of not more than 2 ppm (NMT 2 ppm) for arsenic impurities. A special apparatus called the **Gutzeit apparatus** is used for this test.
    - **Reasons**: Stannous chloride is used to ensure the complete evolution of arsine. Zinc, potassium iodide, and stannous chloride act as powerful reducing agents. Hydrochloric acid makes the solution acidic, which is necessary for the reaction. Lead acetate paper is used to trap any hydrogen sulphide gas that might also be evolved, preventing it from interfering with the arsenic test.
    - **Observation**: The stain in the sample solution should show lesser intensity than that of the standard stain for the sample to pass the limit test.

This concludes our exploration of Unit 1 topics in Pharmaceutical Analysis. You've now gained a foundational understanding of what pharmaceutical analysis is, the techniques used, how to express concentrations, how to prepare and standardise solutions, the types of errors that can occur and how to minimise them, and the vital roles of pharmacopoeias, impurity control, and limit tests. Keep learning, and you'll soon become a true detective in the world of medicines!